BTV-3 Vaccine Comparison

Vaccine	Bluevac-3	Bultavo-3	Syvazul BTV3
Purpose	Active immunisation of sheep to reduce the viraemia, mortality and clinical signs of BTV3. For active immunisation of cattle to reduce the viraemia of BTV3.	Active immunisation of cattle and sheep to prevent clinical signs of BTV3 in both sheep and cattle and prevent viraemia in cattle and reduce viraemia in sheep.	Active immunisation of cattle and sheep to reduce viraemia, mortality and clinical signs of BTV-3
Sheep			
Minimum Age	2 months	1 month	2 months *3 months if DAM vaccinated
Programme	2 doses (3 weeks apart)	1 dose	1 dose
Dose & Route	2 ml subcutaneous	1 ml subcutaneous	2 ml subcutaneous
Cattle			
Minimum Age	2 months	1 month	2 months
Programme	2 doses (3 weeks apart)	2 doses (3 weeks apart)	2 doses (3 weeks apart)
Dose & Route	4 ml subcutaneous	1 ml intramuscular	2 ml intramuscular
General Advice			
Use in Pregnancy	Yes	Yes	Yes
Use in Lactation	Yes	Yes	Yes
Breeding Males	Safety not established	Safety not established	Safety not established
Special Notes	Vaccinate healthy animals only Shake gently before use Once open, avoid repeatedly broaching the vial Change needles regularly	Vaccinate healthy animals only Shake gently before use Warm to 15–25°C before use Avoid bubbles and multiple broaching of the vial Change needles regularly	Vaccinate healthy animals only Shake gently before use Once open avoid repeatedly broaching the vial Change needles regularly
Handling	Keep cool and refrigerate Protect from light	Keep cool and refrigerate Protect from light	Keep cool and refrigerate Protect from light
Mixing with Other Treatments	Do not mix	Do not mix	Do not mix
Shelf Life After Opening	10 hours	10 hours	10 hours