



# GHG Emissions

Laurence Shalloo, Katie Starsmore, Ben Lahart and Jonathan Herron

Animal & Grassland Research and Innovation Centre

Teagasc,

Moorepark,


Fermoy,

Co Cork.

Phone: 025 42 222

web: <http://www.agresearch.teagasc.ie/moorepark/>

Email: [moorepark\\_dairy@teagasc.ie](mailto:moorepark_dairy@teagasc.ie)



**Moorepark2019**



Dairy Research Ireland

funded by Irish Dairy Farmers



Follow us:




# Overview

- Background
- Emissions
- Methane

# Grass fed – Protein efficiency

$$\text{Total Efficiency} = \frac{\text{Proteins produced (whole carcasses, milk)}}{\text{Proteins consumed by livestock (total feed)}}$$

$$\text{Net Efficiency} = \frac{\text{Human edible proteins produced}}{\text{Human edible proteins consumed}}$$



*(adapted from  
Wilkinson, 2011;  
Ertl et al, 2015)*

→ What is human-edible ?

Laisse et al., 2018




# Grass fed – Protein efficiency



Laisse et al., 2018

# Grass fed – Protein efficiency






Grass fed – Environmental Sustainability





# Effect of method and system on GHG emissions



Agricultural Systems 107 (2012) 33–46

Contents lists available at ScienceDirect

Agricultural Systems

journal homepage: [www.elsevier.com/locate/agry](http://www.elsevier.com/locate/agry)



Animal (2012), 6(9), pp. 1512–1527 © The Animal Consortium 2012  
doi:10.1017/S1751731112000316



## Evaluation of the effect of accounting method, IPCC v. LCA, on grass-based and confinement dairy systems' greenhouse gas emissions

D. O'Brien<sup>1,2</sup>, L. Shalloo<sup>1\*</sup>, J. Patton<sup>1</sup>, F. Buckley<sup>1</sup>, C. Grainger<sup>1</sup> and M. Wallace<sup>2</sup>

<sup>1</sup>Livestock Systems Research Department, Animal & Grassland Research and Innovation Centre, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland;

<sup>2</sup>School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Ireland

(Received 15 September 2011; Accepted 12 January 2012; First published online 24 February 2012)


## A life cycle assessment of seasonal grass-based and confinement dairy farms

Donal O'Brien<sup>a,b</sup>, Laurence Shalloo<sup>a,\*</sup>, Joe Patton<sup>a</sup>, Frank Buckley<sup>a</sup>, Chris Grainger<sup>a</sup>, Michael Wallace<sup>b</sup>

<sup>a</sup>Livestock Systems Research Department, Animal & Grassland Research and Innovation Centre, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland

<sup>b</sup>School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Ireland

# Farm profit and carbon footprint of milk



J. Dairy Sci. 98:7394–7407  
<http://dx.doi.org/10.3168/jds.2014-9222>  
 © American Dairy Science Association®, 2015.

## Relating the carbon footprint of milk from Irish dairy farms to economic performance


D. O'Brien,<sup>\*1</sup> T. Hennessy,<sup>†</sup> B. Moran,<sup>†</sup> and L. Shalloo<sup>\*</sup>

<sup>\*</sup>Livestock Systems Research Department, Animal & Grassland Research and Innovation Centre, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland

<sup>†</sup>Rural Economy Research Centre, Teagasc, Athenry, Co. Galway, Ireland



# How does Ireland Compare?



■ On farm (CO<sub>2</sub>e /kgECM) ■ Total (CO<sub>2</sub>e/kgECM)



J. Dairy Sci. 97:1835–1851  
<http://dx.doi.org/10.3168/jds.2013-7174>  
 © American Dairy Science Association®, 2014.

**A case study of the carbon footprint of milk from high-performing confinement and grass-based dairy farms**


D. O'Brien,<sup>\*1</sup> J. L. Capper,<sup>†</sup> P. C. Garnsworthy,<sup>‡</sup> C. Grainger,<sup>\*</sup> and L. Shalloo<sup>\*</sup>

<sup>\*</sup>Livestock Systems Research Department, Animal & Grassland Research and Innovation Centre, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland

<sup>†</sup>Department of Animal Sciences, Washington State University, Pullman 99164

<sup>‡</sup>The University of Nottingham, School of Biosciences, Sutton Bonington Campus, Loughborough, LE12 5RD, United Kingdom


# Emissions per kg milk produced in different EU countries



Source: Evaluation of the livestock sector's contribution to the EU GHG emissions (GGELS) **EC, Joint Research centre, 2010.**

# Carbon footprint of Milk by Country


12





# Carbon footprint of Milk by Country - Errors

12




# Recalculation of the Irish Footprints

- New country specific emission factors have been developed for Ireland by Teagasc research over the past number of years
- Now included in the national GHG inventory compiled by EPA
- New Carbon footprints calculated
  - 1.13 becomes <1.0 kgs CO<sub>2</sub>e per kg FPCM


Emission Factor	Old	New	Reference
Dung kg of N <sub>2</sub> O-N/ kg N excreted	0.02	0.0031	Krol et al. 2016
Urine kg of N <sub>2</sub> O-N/ kg N excreted	0.02	0.0118	Krol et al. 2016
CAN kg of N <sub>2</sub> O-N/ kg N applied	0.01	0.0149	Harty et al. 2016
Urea kg of N <sub>2</sub> O-N/kg N applied	0.01	0.0028	Harty et al. 2016
CAN fertiliser production CO <sub>2</sub> e/kg N	7.11	3.71	Brentrup et al. 2016
Urea fertiliser production CO <sub>2</sub> e/kg N	4.66	3.50	Brentrup et al. 2016

# Carbon footprint of Milk by Country –Recalculation of Irish numbers







## Carbon footprint of Milk by Country – Future protected Urea



# Carbon footprint of Milk by Country – Future protected Urea + lower CP feed<sup>12</sup> (less LUC)




## Carbon footprint of Milk by Country – Future protected Urea + lower CP feed (less LUC)+ performance from grass





# Carbon footprint of Milk by Country – Future Protected Urea + lower CP feed (less LUC)+ performance from grass+ sequestration



# Further strategies

- Mitigation
- Methane
  - Biogenic Methane
    - Metrics GWP100 versus GWP\*
  - Measurement
  - System
  - Additives

# Mitigation strategies

- Footprint
  - Efficiency measures
  - Reduce footprint but could be associated with static or increased absolute emissions (e.g. genetics)
- Absolute emissions
  - Reduce total emissions
  - Footprint?
- Win/Win scenarios reduce footprint and absolute emissions



# Further strategies

- Mitigation
- Methane
  - Biogenic Methane
    - Metrics GWP100 versus GWP\*
  - Measurement
  - System
  - Additives

# Biogenic Methane

- *Biogenic Methane* is emitted from biological processes including livestock.
  - Plants absorb carbon dioxide through the process of photosynthesis
  - Ruminants are then able to break down indigestible cellulose in their rumens
  - carbon that makes up the cellulose is converted to methane
  - After circa 12 years, the methane is converted to carbon dioxide and the cycle starts again.
- In the case of fossil fuel the CO<sub>2</sub> produced is new carbon – Stored for a very long time
- In a situation where methane is constant the same amount of methane that is being produced is being oxidised and therefore there is Little ADDITIONAL warming effect

# Biogenic Methane Metric GWP\*

- Currently all calculations use GWP100
  - Brings everything to 100 year periods
  - Methane has a multiplier of 28
  - Nitrous oxide 265
- Relatively new metric GWP\*
  - Reflects that methane has a half life of 12 years
  - It has a higher multiplier effect at 84

# Biogenic Methane GWP 100


	Livestock*000	Methane (*000 tonnes)	Methane CO2e (*000)
<b>2018</b>	6,594	518.8	12,970
<b>2017</b>	6,674	518.5	12,963
<b>2016</b>	6,613	504.4	12,610
<b>2015</b>	6,422	489.4	12,235
<b>2014</b>	6,243	479.4	11,985
<b>2013</b>	6,309	474.0	11,850
<b>2012</b>	6,253	467.3	11,683
<b>2011</b>	5,925	451.9	11,298
<b>2010</b>	5,918	456.3	11,408
<b>2009</b>	6,232	465.9	11,648
<b>2008</b>	6,304	474.0	11,850
<b>2007</b>	6,248	475.5	11,888
<b>2006</b>	6,340	484.7	12,118
<b>2005</b>	6,390	487.8	12,195
<b>2004</b>	6,212	493.6	12,340
<b>2003</b>	6,223	494.6	12,365
<b>2002</b>	6,333	497.1	12,428
<b>2001</b>	6,408	503.1	12,578
<b>2000</b>	6,330	506.2	12,655
<b>1999</b>	6,558	530.3	13,258
<b>1998</b>	6,952	547.7	13,693




# Biogenic Methane (GWP\*)

	Livestock*000	Methane (*000 tonnes)	Methane CO2e (*000)	Methane GWP* (*000)
<b>2018</b>	6,594	518.8	12,970	2,032
<b>2017</b>	6,674	518.5	12,963	366
<b>2016</b>	6,613	504.4	12,610	- 423
<b>2015</b>	6,422	489.4	12,235	-908
<b>2014</b>	6,243	479.4	11,985	- 555
<b>2013</b>	6,309	474.0	11,850	- 1,076
<b>2012</b>	6,253	467.3	11,683	- 2,056
<b>2011</b>	5,925	451.9	11,298	- 3,690
<b>2010</b>	5,918	456.3	11,408	- 2,799

# Biogenic Methane GWP\* and GWP100



# Cattle numbers December



# Biogenic Methane

- The importance of methane mitigation increases under GWP\* metrics
- Methane effects are magnified
- Significant focus needed on methane
  - System – baseline
  - Genetics
  - Age
  - Additives