

# CROP TRIALS EVENT



***Teagasc Oak Park  
Open Day 2021  
30th June & 1st July***



## Foreword

### Teagasc 2021 Crop Trials Event

Welcome to today's Crop Trials event, which we hope provides you with a welcome opportunity to witness first-hand several aspects of our applied and strategic research programme. While we have had to reduce the size and scope of the event in line with public health measures, through the tour you will hear on-going research related to spring barley, winter wheat, beans, oilseed rape & rye from KT and research staff.

Content includes developing IPM strategies to control BYDV, disease control in spring barley and winter wheat, oilseed rape and rye management, use of liquid N for cereals, establishment systems for beans and field pea, insights from the ASSAP programme as well as grassweed identification and important machinery hygiene actions. The latest spring barley and winter wheat varieties will also be demonstrated by the DAFM variety testing team.

The stakeholder led Crops 2030 strategy published last year was clear in its statement that the future sustainability of the sector is dependent on the need to develop high value markets, while exploiting the authenticity and provenance of Irish material to underpin domestic/export markets. As we expand our programme through national/international partnerships, you will find in Part II of this booklet additional insight into ongoing research being completed across the department.

While attendee numbers have been limited for this year's event we are delighted to be able to host the Farmers Journal live event on the evening of June 30th, which gives an opportunity for all to see the trials and hear research and KT staff discuss the primary issues affecting the tillage sector. Streamed across multiple platforms, this will provide an opportunity for live engagement through an interactive Q&A forum.

In the centre of this booklet you will find a map of the circuit, which has an approximate walking time of 3hr, including stops. At all times, we ask that you adhere to the public health measures in place and the directions of Teagasc staff who are present to assist and ensure a safe and enjoyable day for all.

Hosting the Open day requires significant work and planning, which has been further complicated due to COVID. I therefore wish to thank all Oak Park staff for their commitment and effort in preparing the site, boards and facilities across the campus. In addition, I wish to acknowledge the Teagasc tillage stakeholder group for their support and input to ensure the relevance of our research programme to the industry at large.

Our priority for this event is to maximise socially responsible engagement and research demonstration with attendees and I hope your time here in Oak Park is enjoyable, safe and productive in supporting your business interests. We look forward to inviting you back to a more normal event in 2023.

**Dr. Ewen Mullins**

Head of Crops Research  
Teagasc, Oak Park

---

# Part I


## Boards presented in Crops Trials Event




## Aphid monitoring to support BYDV control

- Understanding impact of climate on aphid movement key for BYDV control
- Knock-down resistance (kdr) in Grain Aphids confers pyrethroid resistance

### Knock down resistance (kdr) in migrating grain




### Suction Tower Locations



Sampling radius of up to 80km per tower

### Aphid Flight



### Impact

- Link aphid migration to forecast (>15C; strong winds and rain negatively affect aphid flight)
- Prevalence of partial insecticide resistance in migrating aphids
- Investigate impact of different migrating aphid species on BYDV spread



Notes:

---



---



---




---



## Accurate BYDV detection


### Why is it important?

- Virus levels vary from field-to-field and across seasons
- Previous systems could only detect if BYDV present/absent in infected tissues
- Critical to be able to quantify BYDV load in both aphids / asymptomatic plants
- Support 'decision-to-spray' actions as part of IPM of BYDV



### How it works

- Exact quantification of virus levels in leaf and aphid samples possible with ddPCR



### Impact

- ddPCR will quantify BYDV load in aphids and a/symptomatic plants
- Virus threshold levels will inform future IPM based spray decisions



Contact details: [Louise.McNamara@Teagasc.ie](mailto:Louise.McNamara@Teagasc.ie)

Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



## Spring barley disease control



Rhynchosporium



Net Blotch

### Fungicide timings?

- Late tillering – protect tillers
- GS49 (awns peeping) – protect green leaf
- Loss of up to 0.4 t/ha if left until GS59 (ear fully out)



Ramularia

### What to use?

- Use mix of actives
- 50% rate of each sufficient
- Azole + SDHI/strobilurin
- Add Folpet for additional Ramularia control



Mildew

Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_




## Managing ramularia



- **Varietal resistance** – don't rely on this
- **Agronomic practises** – don't stress the crop....in the Irish climate??
- **Nutrition** – provide adequate nutrition to minimise stress on the crop
- **Fungicides** – to protect the upper canopy

### Spring barley 2019 – full rates compared



**In absence of CTL careful development of fungicide programme for Ramularia control required**



Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_






## Why break crops?

### Rotations and break crops

- **Yield increase:** after breaks for disease, crop nutrition etc. **+19%** (Knockbeg).
- **Crop diversity:** gives weed control advantages and herbicide options.
- **Improved work distribution:** labour / machinery.
- **More profitable!**



### Knockbeg: Rotation vs continuous wheat



- WOSR, WW, WO, WW, WB
- Performance varied form year to year
- Average benefit: **€208 / ha** for rotation (mean all crops) vs continuous wheat

Notes:

---



---



---



---



## OSR: Growth management

### Approaches

- **Traditional: Fixed N rate**
  - Based on previous crop only
  - 225kg N: 1/3 and 2/3 split applications
- **Canopy Management**
  - Based on post-winter crop green area, soil N and predicted yield
  - Adding N, and timing application, to build canopy for optimum pod number
  - Mainly based on UK research



### Questions

- Are the UK assumptions appropriate?
- Do our climate and soils impact on them?
- Which is better and how do we manage:
  - Early sown, large biomass crops?
  - Later sown, smaller biomass crops?



Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



## OSR – Growth management

### What we did:

- A range of N strategies - canopy management (CM), fixed and zero
- Three biomass levels (sow dates: Mid-Aug, End-Aug, Mid-Sept)
- Defoliated and not (simulating bird damage)
- N content, canopy and yield measurements

### What we found:

- Canopy management principals are largely valid
- Climate and soils have potential for ~50kg/ha
- Scope to be more precise with N, but need estimates of:
  - Soil N
  - Yield potential
- Sowing date impact only if poor establishment conditions
- Defoliation had small effect

### Results snapshot:

	2019		2020	
	N (kg/ha)	Yield (t/ha)	N (kg/ha)	Yield (t/ha)
<b>System</b>				
Fixed	225	<b>4.6</b>	225	<b>4.7</b>
CM 4.5	209	<b>4.3</b>	216	<b>4.7</b>
CM 3.5	149	<b>4.5</b>	156	<b>4.6</b>
Irish CM	-	-	132	<b>5.0</b>

#### Notes:

1. Irish CM assumes extra 50kg/ha of soil N becomes available
2. Data from un-defoliated treatments and mean of 3 sow dates
3. These responses are site and season specific on medium textured soils.

Notes:

---



---



---



---



## Wheat disease control

Management of wheat diseases critical to achieving potential yields



Septoria tritici blotch



Eyespot




Yellow Rust



Fusarium Head Blight



Powdery Mildew



Utilise all components of crop production

### Integrated Pest Management

Crop Selection

•Prevent / limit potential disease establishment

Variety selection

•Know disease resistance strengths and weakness

Crop Agronomy

•Match agronomy to disease pressures / risks (e.g. delayed sowing)

Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



## Yellow rust & septoria control

### Septoria tritici blotch (STB)



- Disease thrives during wet weather
- Typically most damaging disease of Irish wheat crops
- Long latent period can mask asymptomatic disease
- Fungicide resistance increasing and impacting control – anti-resistance strategies essential
- Fungicide timing (final leaf 3 & flag leaf) essential to maximise control

### Yellow Rust

- Increasingly problematic in Irish wheat crops
- Cool & damp conditions favour disease development
- Varietal resistance most effective control measure
- Early identification and control essential to successful management of disease
- Careful selection of active fungicides extremely important



**Varietal selection is critical for STB and Yellow Rust control**

Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_




## Monitoring *Z. tritici* virulence

**Managing and maintaining varietal resistance increasingly vital for successful disease control**



RGT Saki in 2020

- Septoria resistance can be based on minor or major sources of genetic resistance
- Major resistance subject to breakdown as *Z. tritici* can evolve virulence
- In 2019 and 2020 Irish *Z. tritici* evolved virulence to breakdown Cellule- and Cougar-based major resistance



- Illustration indicates virulence to Cougar resistance, detected in Irish *Z. tritici* population in 2020.
- Result indicates reduced potential of several varieties bred from Cougar (C) to resist septoria

Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_


\_\_\_\_\_



## Winter rye management

### Advantages

- High yield potential
- Good disease resistance
- Good nutrient efficiency
- Good drought tolerance
- Spread workload
- High straw yield



### Challenges

- Tall crop – lodging needs to be managed
- Limited market
- Limited pesticide approvals
- Ergot - Low risk in modern hybrids
- Slug damage
- Limited agronomy research



An Roinn Talmhaíochta,  
Bia agus Mara  
Department of Agriculture,  
Food and the Marine

Agronomy research on rye is part funded by DAFM  
Stimulus project DABBING-CAP project no :2019R563

Notes:

---

---

---

---





## Liquid N for cereals

Liquid N = Urea Ammonium Nitrate (UAN)

### Advantages

- Evenness of application,
  - Wide tramlines
  - Less than favourable conditions e.g. windy
- Potentially less overlaps / ins and outs
- Small cost per kg N advantage ?



Liquid N is as efficient as CAN  
in the absence of volatilisation

### Drawbacks

- Urea N can be lost as ammonia
- Can be minimised by including urease inhibitor
- Potential for crop scorch
- On farm storage may be required
- Corrosion

Notes:

---



---



---



---






## Establishment systems for faba beans

### Approach

- Evaluate the impact of cultivation systems on faba bean yield
- 3 seasons x 2 fields (Carlow, Dublin)
- Treatments: Plough + 1 pass (125mm), plough + 1 pass (375mm), plough + Claydon, Claydon into stubble
- Sowing dates: October, February, March



### Future impact:

- Yield comparable with conventional & reduced tillage
- Regardless of the cultivation system used, winter sowing presents higher yield potential

Contact details: [Sheila.Alves@Teagasc.ie](mailto:Sheila.Alves@Teagasc.ie)



An Coimisiún Talmhaíochta,  
Rúnaigh Mairí  
Department of Agriculture,  
Food and the Marine



FACCEP

Notes:

---



---



---




---



## Identifying resistance to chocolate spot in faba beans



- Chocolate spot is the major disease of Irish faba bean crops
- Current varieties lack durable resistance
- EU project (ProFaba) evaluating 200 varieties in Ireland, Spain, France, Germany, Denmark
- Rapid screen to identify resistance to chocolate spot among high yielding lines in ProFaba collection (see below) being developed



**PROFABA**

**W**  
WALSH SCHOLARSHIPS  
RAGB 2019/20



An tAon Talmhaíochta,  
Ní agas Máis  
Department of Agriculture,  
Food and the Marine



**SusCrop - ERA-NET**  
Collaborative Sustainable Crop Production  
FACCEP

Notes:



## Potential of field pea as an alternative protein crop


### Field peas:

- Pea is a high value alternative protein crop
- Good yield potential but no RL for peas
- Lodging issues a brake on expansion


### Approach:

- Varietal trials of winter and spring field peas underway
- Mixed crop systems (peas/beans) for increased pea stand

### Yield potential of spring field pea varieties



### Potential of mixed cropping field pea/faba bean (winter)



### Future impact:

- High yielding spring material confirmed
- Mixed pea/bean cropping has potential to reduce lodging. Further trials underway

Contact details:  
Sheila.Alves@Teagasc.ie





Notes:



## Nitrogen Loss to Water

### Water Framework Directive - WFD

- Nitrates review - open for comment
- Increasing N trend since 2013
  - Drinking water max limit 11.3 mg/l N
  - Estuarine standard below 2.6 mg/l N
- Source of N - 85% from Agriculture
- Soil type, moisture content & N applied
  - South and East of the country at risk



### Tillage systems can leak N

- Slaney & Barrow have highest N loads
  - 27% & 28% tillage
- N loss to water 15 to 65 kg / ha
- Catch Crops, N Efficiency & Excess N
- Big variation year to year - weather
- “Critical Source Areas” & PIP maps



Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



## Tillage vs. Grassland

### Castledockrell - Barley

- Fertiliser N – 140kg/ha
- Organic N - 38 kg N / ha
- 66% tillage
- Free draining soil, slate
- 1020 mm annual rainfall



**Nitrate 7.02mg/l N**  
Target <2.6

### Timoleague - Dairy

- Fertiliser N – 250kg/ha
- Organic N – 170 kg N+
- 85% grassland
- Free draining soil, sandstone
- 1120mm annual rainfall

**Higher N load going on &  
Lower N in the stream ???**



**Nitrate 6.1mg/l N**  
Target <2.6



Notes:

---

---

---

---



# Herbicide-resistant wild oats


## Sensitivity screen

- 12 of 102 populations resistant to one of the ACCase group (DEN, FOP or DIM)
- Low or moderate levels of resistance to DEN (e.g. Axial)
- High levels of resistance to FOP (e.g. Falcon)
- Cross-resistance within field

## Solutions

- No ALS resistance to date
- Use ALS herbicides with cultural control; *Integrated Weed Management* (IWM), to avoid resistance

Effective dose rate required to kill 90% of the treated plants for resistant (red) and susceptible (green) populations to:



**\*Blue line denotes label rate**




Notes: \_\_\_\_\_

\_\_\_\_\_


\_\_\_\_\_

\_\_\_\_\_

# Teagasc 2021 Crop Trials Event







Coronavirus  
**COVID-19**  
Public Health  
Advice

# Coronavirus **COVID-19**

**Stay safe.  
Protect  
each other.**




## **Wash**

your hands well  
and often to avoid  
contamination.



## **Cover**

your mouth and nose  
with a tissue or sleeve  
when coughing or  
sneezing and discard  
used tissue safely



## **Distance**

yourself at least  
2 metres (6 feet) away  
from other people,  
especially those who  
might be unwell

**#holdfirm**

## **For more information**

[www.gov.ie/health-covid-19](http://www.gov.ie/health-covid-19)  
[www.hse.ie](http://www.hse.ie)





## Herbicide-resistant black-grass



### Sensitivity screen

- 8 of 18 populations resistant
- Resistance (multiple) to both ALS and ACCase herbicides found in 5 populations
- Cross-resistance to ACCase-only herbicides found in 3 populations

### Solutions

- For ACCase-only resistant populations, use ALS and IWM.
- For multiple-resistant populations, cultural IWM, including grass/leys, for a minimum of 5 years needed to eliminate seedbank

Effective dose rate required to kill 90% of the treated plants for resistant (red) and susceptible (green) populations to:



\*Blue line denotes label rate



Ireland's European Structural and Investment Funds Programmes 2014-2020  
Co-funded by the Irish Government and the European Union



An Roinn Talmhaíochta, Bia agus Mara  
Department of Agriculture, Food and the Marine



The European Agricultural Fund for Rural Development  
Investing in rural areas

Notes:

---



---




---



---



## Integrated strategies for managing grass weeds



Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_




## Integrated weed management (IWM) Strategies for blackgrass control

### ECT Focus Farm Waterford

- Blackgrass 1<sup>st</sup> appeared in 2012
- By 2017 winter cropping had become unviable
- Blackgrass population on the farm – **RESISTANT** to both ALS and ACCase chemistry

### IWM Strategy used on the farm for last 3 years

- Stale seedbeds
- Spring Cropping
- Increased seeding rates
- Good on-farm Hygiene has confined Blackgrass to this area



Blackgrass plant counts	2019	2020
Range (plants/m <sup>2</sup> )	0 - 142	1 - 107
Average (plants/m <sup>2</sup> )	21	11
Average reduction	-	52%

**Problem still exists and will influence all decisions for at least next 5 years**



Notes:

---



---



---



---



## Stop grass weed spread

### Machine cleaning: Why?

- To stop the introduction and spread of problematic weeds. One black-grass plant = 6000 viable seeds.
- To keep crop production viable – key IWM measure



### Which machines and when?

- All in crop contact: baler, combine, rake, tractors, trailers etc.
- Whether owned, contractor or other.
- When moving from any farm or suspect area on own farm

### Resources

- Long handled brushes, leaf blower, compressed air and long lance, torch, all openings accessible safely.
- PPE and face mask. Safe working practice.
- Knowledge re safe cleaning routine for specific machine
- All involved aware of the need for cleaning – critical.

### Known black-grass: Don't harvest or bale!

- Destroy, ensile or hand rogue and half-day combine clean!

Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



## Cleaning practice

### All machines

- Before season start: clean, no oil leaks and guards in place.
- Run power-driven machines before and after each cleaning.
- Clean before leaving field and quickly again after moving.

### Balers

- Eject part-bale before cleaning and leaving field.
- Use leaf blower / brush to clean all areas including pick-up, bale chamber, chassis, knotter/netter and under guards.
- Check tractor underbody.
- Note field/position where first bale is made for later checking.

### Combines

- Clean header thoroughly and run complete machine while varying/opening all fan and sieve adjustments.
- Clean all external surfaces and all accessible internal areas.
- Open all elevator covers and run combine (no bystanders).
- Following move, quick clean and run combine (stationary)
- Note field and pathway of first work on farm for later checking.

Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

# Part II

**Snapshot of  
additional research  
currently underway**



### Fungicide sensitivity of Irish *Pyrenopeziza brassicae* populations

#### Background:

- Light leaf spot caused by *P. brassicae* is a major disease of Irish oilseed rape crops
- Limited information known on Irish populations

#### Approach:

- Detailed surveys of Irish population conducted during 2019 and 2020
- Fungicide sensitivity and population analysis completed

#### Outputs to date:

- Irish population remains sensitive to SDHI and QoI fungicides
- Varying levels of resistance detected to azole fungicides with key alterations detected



>500 isolates of *P. brassicae* isolated from Irish OSR crops in 2019 and 2020

#### Future impact:

- Detailed population analysis ongoing to determine genetic structure of Irish population
- Molecular assays being developed to rapidly screen future populations

#### Contact:

- Diana Bucur, CELUP email: [diana.bucur@teagasc.ie](mailto:diana.bucur@teagasc.ie)






## Biostimulants for sustainable wheat production

### Background:


- The loss of chemical efficacy and increased EU legislation around PPP use is challenging the sustainability of crops such as wheat to offset losses against *Zymoseptoria tritici*
- Studies indicate that the soil-borne bacterium *Ensifer adhaerens* has the capacity to release nutrients (e.g. Fe, Si, Al via soil mineralisation) and has antifungal properties

### Outputs:

- Treating seed with *Ensifer adhaerens* OV14 improves wheat seedlings vigour index by an average of 23% (cv. Lilli) and 40% (cv. Rockefeller) compared to untreated seed



- When co-cultivated under *in-vitro* conditions, *Ensifer adhaerens* OV14 shows the ability to inhibit *Z. tritici* growth by up to 80%



### Future Impacts:

- Studying more crop species of high importance in Ireland will contribute to fully characterise *Ensifer adhaerens* OV14 as a new biostimulant agent able to decrease chemical fungicides and fertilizers input



Contact details: [Elena.Grosu@Teagasc.ie](mailto:Elena.Grosu@Teagasc.ie)





# Potential of distilling wheat

## Background:

- Significant imports of maize to produce whiskey
- Wheat is used in whiskey production abroad
- Can Irish wheat be used in Irish distilleries?

## Approach:

- Integrated approach
- Agronomy work examining suitability of distilling varieties and identifying suitable fertiliser strategies
- Lab-scale analysis of suitability of wheat (and other grains) for use in Irish distilleries
- Engagement with drinks sector to highlight results and ensure work is relevant to industry

**Outputs to date:** The project is in its first year

## Future impact:

- Management guidelines for the production of distilling wheat
- Laboratory capability for testing new varieties for their suitability for Irish distilling

Contact details: [john.spink@teagasc.ie](mailto:john.spink@teagasc.ie)  
[richie.hackett@teagasc.ie](mailto:richie.hackett@teagasc.ie)

This research is funded by DAFM Stimulus project DABBING-  
CAP project no :2019R563



An Roinn Talmhaíochta,  
Bia agus Mara  
Department of Agriculture,  
Food and the Marine





# Leguminous cover crops

- Background:**
- Cover crops accumulate soil N overwinter
  - Non-leguminous cover crops generally provide little N to subsequent cereal
  - Leguminous cover crops can fix atmospheric N
  - Can leguminous cover crops supply significant amounts of N to the cereal?

- Approach:**
- Grow a range of leguminous cover crops (vetches, peas, lentil, clovers)
  - Sow spring barley after incorporation of the cover crops
  - Apply a range of fertiliser N levels to barley
  - Compare yield (and protein) with and without leguminous cover crop

**Outputs to date:** Data on spring barley grain yield and protein content with and without leguminous cover crop

**Future impact:** Guidance on use of leguminous cover crops in spring cereal systems

Contact details: [richie.hackett@teagasc.ie](mailto:richie.hackett@teagasc.ie)





## Generating practical advice for leatherjacket control

### Background:

Leatherjackets (LJ) are the larvae of the crane fly, & are major agricultural pests in grassland and cereals.



Key chemical control (chlorpyrifos) prohibited by the EU in 2019 – what next?

### Surveyed farm sites 2018-2021



### Outputs:


- Survey of 123 fields to identify main species completed
- Identified key factors that hinder larval development
- 2018/19 grassland survey suggested a correlation between pH, P, Mg and LJ density
- Profiling soil microbes to identify species with potential to suppress larvae

### Future Impact:

- Identify risk factors (soil characteristics, microbial communities associated with high LJ densities)
- Enhance understanding of LJ biology to allow development of targeted control approaches


### Species ID results 2018/2019

#### Grassland



n larvae = 217  
n fields = 30

#### Tillage



n larvae = 141  
n fields = 31

\*2018/2019 Irish samples

Contact: [colin.mccormack@teagasc.ie](mailto:colin.mccormack@teagasc.ie)

WALSH SCHOLARSHIPS  
PROGRAMME





# Unlocking the aphid genome

## Background:

- The grain aphid is a major agricultural pest of cereals and vector of BYDV
- A clone with partial resistance to insecticides has emerged in recent years
- The genome is like a roadmap with the information needed to build and maintain an organism. Until now, the sequence of DNA units making up the aphid genome was unknown; hindering our understanding of the molecular mechanisms of insecticide resistance

## Outputs:

- We have produced the first ever assembly of the grain aphid's genome and identified the sequence of its over 400 million units of DNA
- We have also identified and characterised over 31,000 genes in the grain aphid genome



## Future Impacts:

- The genome of a clone with partial resistance to insecticides will facilitate research into molecular mechanisms of insecticide resistance, and facilitate development of improved diagnostic tools

Contact details: [Stephen.byrne@Teagasc.ie](mailto:Stephen.byrne@Teagasc.ie)





# White Clover Genomics

## Background:

- White clover can contribute to both the economic and environmental sustainability of pastoral based production systems
- Opportunities exist to accelerate genetic gain in white clover breeding using technologies such as genomic selection

## Outputs:

- We have established DNA profiles on breeding populations
- We have used these DNA profiles with field evaluations and developed promising genomic prediction models for seasonal forage yield



## Future Impacts:

- It is expected that we can accelerated development of white clover cultivars with improved forage yield using the models developed at Oak Park
- Genomic selection will accelerate genetic gain for key traits and help increase on-farm profitability through reduced inputs and increased productivity

Contact details: [katie.hetherington@Teagasc.ie](mailto:katie.hetherington@Teagasc.ie)






## Improving pasture digestibility

### Background:

- While there have been clear gains in breeding for dry matter yield, improvement in forage digestibility has lagged behind
- Genomic Selection (GS) has the potential to greatly contribute to improving complex forage quality traits



### Outputs:

- We have developed promising genomic prediction models for forage quality parameters such as digestibility

### Future Impacts:

- GS can enable a selection cycle to be completed in a single year (compared to seven with field evaluations), enable more plants be evaluated, and improve accuracy of selection
- Improving forage digestibility will lead to increased animal performance

Contact details: [Agnieszka.Konkolewska@Teagasc.ie](mailto:Agnieszka.Konkolewska@Teagasc.ie)



An Roinn Talmhaíochta,  
Bia agus Mara  
Department of Agriculture,  
Food and the Marine



## Optimal faba beans management

- Background :**
- Yield instability remains a primary factor challenging greater faba bean uptake
  - Large scale trials were completed to identify optimal management practises to maximise yield potential

Parameter	Optimal practise/conditions to support yield
<b>Sowing window</b>	Winter beans: October Spring Beans: Mid-February to mid-March
<b>Seed rate</b>	Winter beans: < 30 seeds m <sup>-2</sup> Spring beans: 40 seeds m <sup>-2</sup>
<b>Soil nutrition</b>	6 < pH < 7 P-fertilisation in low P index soils can increase yield up to 40%
<b>Cultivation system</b>	Conventional and reduced tillage systems give comparable yields
<b>Disease control</b>	Chocolate spot (major), downy mildew, rust Application: 2x from flowering. No additional benefit of 3 <sup>rd</sup> application based on trials at Oak Park/Kildalton. Further trials planned at additional locations
<b>Pests</b>	Evidence of leaf notching needs to be addressed with insecticide application to minimise impact of pea & bean weevil
<b>Dry spells</b>	Early sowing will render a crop less susceptible to dry spells between May-July
<b>Harvest</b>	Mid September to October (small % of green stems remaining) Aim for < 20% moisture content



**Future impact :** Adoption of measures will minimise yield instability and support economic returns

Contact details: [Sheila.Alves@Teagasc.ie](mailto:Sheila.Alves@Teagasc.ie)



An Roinn Talmhaíochta,  
Bia agus Mara  
Department of Agriculture,  
Food and the Marine















## U-Protein: adding value to protein crops

- Background:**
- Goal of U-Protein is to unlock the added value potential of protein crops
  - Expand the alternative use and circularity potential of crops via novel post-harvest processing / formulation

- Approach:**
- Evaluate protein crops with potential under Irish conditions to deliver high protein content
  - Tailor the grain amino acid profile via agronomy measures to support high value protein processing be that for food/animal feed outputs

### Protein content

25%	29%	20%	20%	23%	22%	30%	25%	40%	20%
									
Dry beans	Lentils	Cowpeas	Pigeon peas	Dry peas	Chickpeas	Faba beans	Bambara beans	Lupins	Vetches

- Outputs:**
- Project commenced earlier this year and initial results will be presented at 2022 Tillage Conference

- Future impact:**
- Identify alternative crops for future rotations that deliver high value feedstock for food/feed processing

Contact details:  
[lucie.cases@teagasc.ie](mailto:lucie.cases@teagasc.ie)







## Barley Against Rhynchosporium Disease


### Background:

- Rhynchosporium is a major barley disease, reducing tiller survival and yield
- Sequencing technology allows us to examine the genetic response of different barley cultivars to Rhynchosporium
- This allows us to identify genes linked with varietal resistance or contributing to susceptibility, which can be important targets for breeding programmes

### Outputs:

- Barley responds diversely to different Rhynchosporium strains
- Barley varieties Cassia and Infinity have an overlapping 'supercore' set of 983 genes that are activated in response to infection
- We have identified a core number of genes that promote susceptibility to Rhynchosporium

Barley Variety	Resistance rating	Number of Core response genes
KWS Cassia	4	1846
KWS Infinity	7	1717



Rhynchosporium infection on var Cassia

### Future Impacts:

Understanding the genetics behind infection will allow us to identify genes associated with resistance, which will be targets for the breeding of barley varieties with more durable resistance.



Contact details: [tara.oconnor@Teagasc.ie](mailto:tara.oconnor@Teagasc.ie)



Notes:

[illegible]

